

Harmonic Analysis on Real Reductive Symmetric Spaces

Patrick Delorme*

Abstract

Let G be a reductive group in the Harish-Chandra class e.g. a connected semisimple Lie group with finite center, or the group of real points of a connected reductive algebraic group defined over $\mathbb{R}$. Let σ be an involution of the Lie group G , H an open subgroup of the subgroup of fixed points of σ . One decomposes the elements of $L^2(G/H)$ with the help of joint eigenfunctions under the algebra of left invariant differential operators under G on G/H .

2000 Mathematics subject classification: 22E46, 22F30, 22E30, 22E50, 33C67.

Keywords and Phrases: Reductive symmetric space, Plancherel formula, Meromorphic continuation of Eisenstein integrals, Temperedness, Truncation, Maass-Selberg relations.

1. Introduction

Let G be a real reductive group in the Harish-Chandra class [H-C1], e.g. a connected semisimple Lie group with finite center, or the group of real points of a connected reductive algebraic group defined over $\mathbb{R}$. Let σ be an involution of the Lie group G , H an open subgroup of the subgroup of fixed points of σ .

Important problems of harmonic analysis on the so-called reductive symmetric space G/H are :

(a) to make the simultaneous spectral decomposition of the elements of the algebra $\mathbb{D}(G/H)$ of left invariant differential operators under G on G/H . In other words, one wants to write the elements of $L^2(G/H)$ with the help of joint eigenfunctions under $\mathbb{D}(G/H)$.

(b) to decompose the left regular representation of G in $L^2(G/H)$ into an Hilbert integral of irreducible unitary representations of G : this is essentially the Plancherel formula.

*Institut de Mathématiques de Luminy, U.P.R. 9016 du C.N.R.S., Faculté des Sciences de Luminy, 163 Avenue de Luminy, Case 930, 13288 Marseille Cedex 09, France. E-mail: delorme@iml.univ-mrs.fr

(c) to decompose the Dirac measure at eH , where e is the neutral element of G , into an integral of H -fixed distribution vectors : this is essentially the Fourier inversion formula.

These problems were solved for the “group case” (i.e. the group viewed as a symmetric space : $G = G_1 \times G_1$, $\sigma(x, y) = (y, x)$, H is the diagonal of $G_1 \times G_1$) by Harish-Chandra in 1970s (see [H-C1,2, 3]), the Riemannian case (H maximal compact) had been treated before (see [He]). Later, there were deep results by T. Oshima [O1]. When G is a complex group and H is a real form, the Problems (a), (b), (c) were solved by P. Harinck, together with an inversion formula for orbital integrals ([Ha], see also [D3] for the link of her work with the work of A. Bouaziz on real reductive groups).

Then, E. van den Ban and H. Schlichtkrull, on the one hand, and I, on the other hand, obtained different solutions to problems (a), (b), (c). Moreover, they obtained a Paley-Wiener theorem (see [BS3] for a presentation of their work). I present here my point of view, with an emphasize on problem (a), because it simplifies the formulations of the results (nevertheless, the important aspect of representation theory is hidden). It includes several joint works, mainly with J. Carmona, and also with E. van den Ban and J.L. Brylinski. Several works of T. Oshima, linked to the the Flensted-Jensen duality, alone and with T. Matsuki are very important in my proof, as well as earlier results of E. van den Ban and H. Schlichtkrull.

I have to acknowledge the deep influence of Harish-Chandra’s work. The crucial role played by the work [Be] of J. Bernstein on the support of the Plancherel measure, and some part of Arthur’s article on the local trace formula [A] will be appearant in the main body of the article.

2. Temperedness of the spectrum

Let θ be a Cartan involution of G commuting with σ , let K be the fixed point set of θ . Let $\mathfrak{g}$ be the Lie algebra of G , etc. Let $\mathfrak{s}$ (resp. $\mathfrak{q}$) be the space of elements in $\mathfrak{g}$ which are antiinvariant under the differential of θ (resp. σ). Let $\mathfrak{a}_\theta$ be a maximal abelian subspace of $\mathfrak{s} \cap \mathfrak{q}$. If P is a $\sigma\theta$ -stable parabolic subgroup of G , containing $A_\theta := \exp \mathfrak{a}_\theta$, we denote by $P = M_P A_P N_P$ its Langlands σ -decomposition. More precisely A_P is the subgroup of the elements a of the split component of the Levi factor $L_P = P \cap \theta(P)$ such that $\sigma(a) = a^{-1}$. Here M_P is larger than that for the usual Langlands decomposition.

In order to simplify the exposition we will make the following simplifying assumption:

Hypothesis: *For any P as above, HP is the unique open (H, P) -double coset.*

When $\sigma = \theta$ (the case of a riemannian symmetric space) or the “group case”, this hypothesis is satisfied.

To get the Plancherel formula, it is useful to use K -finite functions. They are often replaced by τ -spherical functions. Here (τ, V_τ) is a finite dimensional unitary representation of K and a τ -spherical function on G/H is a function $f : G/H \rightarrow V_\tau$ such that $f(kx) = \tau(k)f(x)$, $k \in K$, $x \in G/H$.

Some spaces of τ -spherical functions on G/H play a crucial role in the theory, namely:

(a) $\mathcal{C}(G/H, \tau)$: the Schwartz space of τ -spherical functions on G/H which are rapidly decreasing as well as their derivatives by elements of the enveloping algebra $U(\mathfrak{g})$ of $\mathfrak{g}$ (see [B2]).

(b) $\mathcal{A}(G/H, \tau)$: the space of smooth τ -spherical functions on G/H which are $\mathbb{D}(G/H)$ finite. Here $\mathcal{A}$ is used to evoke automorphic forms.

(c) $\mathcal{A}_{temp}(G/H, \tau)$: the space of elements of $\mathcal{A}(G/H, \tau)$ which have tempered growth as well as their derivatives by elements of $U(\mathfrak{g})$ ([D2]). Integration of functions on G/H defines a pairing between $\mathcal{A}_{temp}(G/H, \tau)$ and $\mathcal{C}(G/H, \tau)$.

(d) $\mathcal{A}_2(G/H, \tau)$: the space of square integrable elements of $\mathcal{A}(G/H, \tau)$. This is a subspace of the three preceding spaces.

One has:

Theorem 1 ([D2]): *The space $\mathcal{A}_2(G/H, \tau)$ is finite dimensional.*

This is deduced from the theory of discrete series for G/H initiated by M. Flensted-Jensen [F-J] and achieved by T. Oshima and T. Matsuki, using the Flensted-Jensen duality [OM]. One has also to use the behaviour of the discrete series under certain translation functors, studied by D. Vogan [V] and a result of H. Schlichtkrull [S] on the minimal K -types of certain discrete series.

The next result follows from the work of J. Bernstein [Be] on the support of the Plancherel measure.

Theorem 2 ([CD1], Appendice C): *Every function in $\mathcal{C}(G/H, \tau)$ can be canonically desintegrated as an integral of elements of $\mathcal{A}_{temp}(G/H, \tau)$.*

This information appeared to be crucial at the end of our proof.

3. The continuous spectrum: Eisenstein integrals

Let $P = MAN$ the Langlands σ -decomposition of a $\sigma\theta$ -stable parabolic subgroup P of G . Let ρ_P be the half sum of the roots of $\mathfrak{a}$ in $\mathfrak{n}$ and $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$ be such that the real part of $\lambda - \rho_P$ is strictly dominant with respect to the roots of $\mathfrak{a}$ in $\mathfrak{n}$. Let τ_M be the restriction of τ to $M \cap K$. Then, if $x \in G/H$ and $\psi \in \mathcal{A}_2(M/M \cap H, \tau_M)$, the following integral is convergent:

$$E(P, \psi, \lambda)(x) := \int_K \tau(k^{-1}) \Psi_{\lambda}(kx) dk,$$

where $\Psi_{\lambda}(x) = 0$ if $x \notin PH$, and $\Psi_{\lambda}(x) = a^{-\lambda + \rho_P} \psi(m)$ if $x = namH$ with $n \in N$, $a \in A$, $m \in M$. Moreover $E(P, \psi, \lambda)$ is an element of $\mathcal{A}(G/H, \tau)$. Eisenstein integrals are the τ -spherical versions of K -finite functions of the form: $gH \mapsto \langle \pi'(g)\xi, v \rangle$, where π' is the contragredient of a generalized principal series π , ξ is a certain H -fixed distribution vector of π , v is a K -finite vector of π .

Theorem 3 ([BrD]): *The function $\lambda \mapsto E(P, \psi, \lambda)$ admits a meromorphic continuation in $\lambda \in \mathfrak{a}_{\mathbb{C}}^*$. This meromorphic continuation, denoted in the same way, multiplied by a suitable product, p_{ψ} , of functions of type $\lambda \mapsto (\alpha, \lambda) + c$, where α is a root of $\mathfrak{a}$ and $c \in \mathbb{C}$, is holomorphic around $i\mathfrak{a}^*$.*

This meromorphic continuation is an interesting feature of the theory. For the “group case”, it comes down to the meromorphic continuation of Knapp-Stein intertwining integrals. My proof with Brylinski uses D -modules arguments.

The case where P is minimal had been treated separately by E. van den Ban [B1] and G. Olafsson [Ol]. One has also to mention the earlier work of T. Oshima and J. Sekiguchi [OSe] on the spaces of type G/K_ε .

The proof which gives the best results uses a method of tensoring by finite dimensional representations. It is a joint work with J. Carmona. It was initiated by D. Vogan and N. Wallach (see [W], chapter 10) for the meromorphic continuation of the Knapp-Stein intertwining integrals. For symmetric spaces and the most continuous spectrum, the proof is due to E. van den Ban [B2]. This proof uses Bruhat’s thesis and tensoring by finite dimensional modules. This implies rough estimates for Eisenstein integrals, which generalize those obtained by E. van den Ban when P is minimal [B2].

Theorem 4 ([D1]): *If $\lambda \in i\mathfrak{a}^*$ is such that $E(P, \psi, \lambda)$ is well defined, then it is tempered, i.e. is an element of $\mathcal{A}_{temp}(G/H, \tau)$.*

This is a natural result but the proof is quite long. It uses the behaviour under translation functors of H -fixed distribution vectors of discrete series and of generalized principal series, and also of the Poisson transform. Moreover the duality of M. Flensted-Jensen, [F-J], and a criteria of temperedness due to Oshima [O2] play a crucial role (apparently, J. Carmona has a way to avoid boundary values).

With the help of this theorem and by using techniques due to E. van den Ban [B2], the rough estimates for Eisenstein integrals can be improved to get uniform sharp estimates for $p_\psi(\lambda)E(P, \psi, \lambda)$, $\lambda \in i\mathfrak{a}^*$ (cf. [D1]).

4. C -functions

Let P be as above and let L be equal to MA . The theory of the constant term, due to J. Carmona [C1] (Harish-Chandra for the group case, [H-C1]), gives a linear map from $\mathcal{A}_{temp}(G/H, \tau)$ into $\mathcal{A}_{temp}(L/L \cap H, \tau_L)$, $\varphi \mapsto \varphi_P$, characterized by :

$$\lim_{t \rightarrow +\infty} \delta_P^{1/2}((\exp tX)l)\varphi((\exp tX)l) - \varphi_P((\exp tX)l) = 0,$$

where $l \in L/L \cap H$, $X \in \mathfrak{a}_P$ is P -dominant and δ_P is the modular function of P .

Let Q be a $\sigma\theta$ -stable parabolic subgroup of G with the same θ -stable Levi subgroup L other than P . Let $W(\mathfrak{a})$ be the group of automorphisms of $\mathfrak{a}$ induced by an element of $Ad(G)$. One defines meromorphic functions on $\mathfrak{a}_{\mathbb{C}}^*$, $\lambda \mapsto C_{Q|P}(s, \lambda)$ with values in $End(\mathcal{A}_2(M/M \cap H, \tau_M))$ such that :

$$E(P, \psi, \lambda)_Q(ma) = \sum_{s \in W(\mathfrak{a})} (C_{Q|P}(s, \lambda)\psi)(m)a^{-s\lambda}, \quad m \in M, a \in A, \lambda \in i\mathfrak{a}^*,$$

or rather for λ in an open dense subset of $i\mathfrak{a}^*$.

The C -functions allow to normalize Eisenstein integrals as follows:

$$E^0(P, \psi, \lambda) := E(P, C_{P|P}(1, \lambda)^{-1}\psi, \lambda).$$

5. Truncation, Maass-Selberg relations and the regularity of normalized Eisenstein integrals

Let P be as above and let $P' = M'A'N'$ be the Langlands σ -decomposition of another $\sigma\theta$ -stable parabolic subgroup of G . Let ψ (resp. ψ') be an element of $\mathcal{A}_2(M/M \cap H, \tau_M)$ (resp. $\mathcal{A}_2(M'/M' \cap H, \tau_{M'})$). One chooses p_ψ as in Theorem 3, such that the product of p_ψ with the C -functions are holomorphic in a neighbourhood of $i\mathfrak{a}^*$ which is a product of $i\mathfrak{a}^*$ with a neighbourhood of 0 in $\mathfrak{a}^*$. We do the same for ψ' . One defines $F(\lambda) := p_\psi(\lambda)E(P, \psi, \lambda)$. One defines similarly F' .

One assumes, for the rest of the article, that G is semisimple. This is just to simplify the exposition. One chooses $T \in \mathfrak{a}_\theta$, regular with respect to the roots of $\mathfrak{a}_\theta$ in $\mathfrak{g}$. Let C_T^1 be the convex hull of $W(\mathfrak{a}_\theta)T$ and let C_T be equal to the subset $K(\exp C_T^1)H$ of G/H .

Theorem 5 ([D2]):

(i) *One gets an explicit expression $\omega^T(\lambda, \lambda')$, involving the C -functions (see an example below) and vanishing when A and A' are not conjugate under K , which is asymptotic to*

$$\Omega^T(\lambda, \lambda') := \int_{C_T} (F(\lambda)(x), F'(\lambda')(x)) dx,$$

when T goes to infinity and $\lambda \in i\mathfrak{a}^*$, $\lambda' \in i\mathfrak{a}'^*$. More precisely for $\delta > 0$ there exists $C > 0$, $k \in \mathbb{N}$ and $\varepsilon > 0$, such that for all T satisfying $\|\alpha(T)\| \geq \delta \|T\|$ for every root α of $\mathfrak{a}_\theta$ in $\mathfrak{g}$, one has:

$$\|\Omega^T(\lambda, \lambda') - \omega^T(\lambda, \lambda')\| \leq C(1 + \|\lambda\|)^k (1 + \|\lambda'\|)^k e^{-\varepsilon\|T\|}.$$

(ii) *Moreover ω^T is analytic in $(\lambda, \lambda') \in i\mathfrak{a}^* \times i\mathfrak{a}'^*$.*

This generalizes a result of J. Arthur for the group case [A], Theorem 8.1. My proof is quite similar, but I was able to avoid his use of the Plancherel formula.

(ii) is an easy consequence of (i). In fact, the explicit form of ω^T implies that it is meromorphic around $i\mathfrak{a}^* \times i\mathfrak{a}'^*$. Moreover Ω^T is holomorphic, hence locally bounded, around $i\mathfrak{a}^* \times i\mathfrak{a}'^*$. From the inequality in (i), one deduces that ω^T is locally bounded, hence holomorphic, around $i\mathfrak{a}^* \times i\mathfrak{a}'^*$.

We will now show, by an example, how the explicit form of ω^T and its analyticity in $(\lambda, \lambda') \in i\mathfrak{a}^* \times i\mathfrak{a}'^*$ imply the Maass-Selberg relations.

Let σ be equal to θ , $H = K$, and τ be the trivial representation. Let P, P' be minimal parabolic subgroups of G . Then $\dim \mathcal{A}_2(M/M \cap H, \tau) = 1$ and the C -functions are scalar valued. One assumes $\mathfrak{g}$ to be semisimple and that the dimension of A is one. Then $W(\mathfrak{a})$ has 2 elements, ± 1 , and one has the following explicit expression of ω^T :

$$\omega^T(\lambda, \lambda') = p_\psi(\lambda)p_{\psi'}(\lambda') \sum_{s=\pm 1, s'=\pm 1} e^{s\lambda T - s'\lambda' T} C_{P|P}(s, \lambda) \overline{C_{P|P}(s', \lambda')} (s\lambda - s'\lambda')^{-1}.$$

Thus $\omega^T(\lambda, \lambda')$ is the sum of a product of $(\lambda - \lambda')^{-1}$ by an analytic function with a product of $(\lambda + \lambda')^{-1}$ by an analytic function. The analyticity at (λ, λ) implies

easily that the factor in front of $(\lambda - \lambda')^{-1}$ vanishes for $\lambda = \lambda'$. Hence we get $|C_{P|P}(1, \lambda)|^2 = |C_{P|P}(-1, \lambda)|^2$, $\lambda \in i\mathfrak{a}^*$. This is one of the Maass-Selberg relations (cf. [D2], Theorem 2, and the work with J. Carmona [CD2], Theorem 2 for the general case, see [B1], [B2] for the case where P is minimal). These relations imply that the C -functions attached to normalized Eisenstein integrals are unitary, when defined, for λ purely imaginary. Hence they are locally bounded. This implies that they are holomorphic around the imaginary axis. This implies in particular some holomorphy property of the constant term of normalized Eisenstein integrals. From this, with the help of [BCD], one deduces:

Theorem 6 (Regularity theorem for normalized Eisenstein integrals, [CD2], [BS1] for P minimal): *The normalized Eisenstein integrals are holomorphic in a neighbourhood of the imaginary axis.*

6. Fourier transform and wave packets

Theorem 7 ([CD2], [BS1] for P minimal): *For $f \in \mathcal{C}(G/H, \tau)$, one has $\mathcal{F}_P^0 f \in \mathcal{S}(i\mathfrak{a}^*) \otimes \mathcal{A}_2(M/M \cap H, \tau_M)$, where $\mathcal{F}_P^0 f$ is characterized by:*

$$((\mathcal{F}_P^0 f)(\lambda), \psi) = \int_{G/H} (f(x), E^0(P, \psi, \lambda)(x)) dx, \quad \lambda \in i\mathfrak{a}^*, \quad \psi \in \mathcal{A}_2(M/M \cap H, \tau_M),$$

here $\mathcal{S}(i\mathfrak{a}^*)$ is the usual Schwartz space.

This theorem follows from the sharp estimates of Eisenstein integrals.

Theorem 8 ([BCD]): *If Ψ is an element of $\mathcal{S}(i\mathfrak{a}^*) \otimes \mathcal{A}_2(M/M \cap H, \tau_M)$, one has $\mathcal{J}_P^0 \in \mathcal{C}(G/H, \tau)$, where :*

$$\mathcal{J}_P^0(x) := \int_{i\mathfrak{a}^*} E^0(P, \Psi(\lambda), \lambda) d\lambda, \quad x \in G/H.$$

This theorem follows from the regularity theorem and from the joint work with E. van den Ban and J. Carmona, [BCD].

Now we want to compute $\mathcal{F}_P^0 \mathcal{J}_P^0$. For this purpose one has to study the integral:

$$I := \int_{G/H} \left(\int_{i\mathfrak{a}^*} \alpha(\lambda) E^0(P, \psi, \lambda)(x) d\lambda, E^0(P', \psi', \lambda')(x) \right) dx.$$

One truncates the integral on G/H to C_T and let T goes to infinity (far from the walls). Let us denote the truncated inner product of the normalized Eisenstein integrals by ${}^0\Omega^T(\lambda, \lambda')$. Using Fubini's theorem one has:

$$I = \lim_{T \rightarrow \infty} \int_{i\mathfrak{a}^*} \alpha(\lambda) {}^0\Omega^T(\lambda, \lambda') d\lambda.$$

As for unnormalized Eisenstein integrals, one has an asymptotic evaluation of ${}^0\Omega^T(\lambda, \lambda')$ by an explicit expression ${}^0\omega^T(\lambda, \lambda')$. One can replace ${}^0\Omega^T$ by ${}^0\omega^T$ in the previous formula. By using an expression of ${}^0\omega^T(\lambda, \lambda')$, viewed as a distribution in λ , for fixed λ' , involving Fourier transforms of cones ([D2], Theorem 3) one gets:

Theorem 9 ([CD2]): *Let $\mathbb{F}$ be a set of representative of σ -association classes of $\sigma\theta$ -stable parabolic subgroups. Here σ -association means that the $\mathfrak{a}$ are conjugated under K . Define:*

$$\mathcal{P}_\tau = \sum_{P \in \mathbb{F}} ((W(\mathfrak{a}_P))^{-1} \mathcal{I}_P^0 \mathcal{F}_P^0.$$

Then $\mathcal{P}_\tau$ is an orthogonal projection operator in $\mathcal{C}(G/H, \tau)$ endowed with the L^2 scalar product.

7. The Plancherel formula

Essentially, the solution to problem (a) is contained in the following:

Theorem 10 ([D4]): *The projection $\mathcal{P}_\tau$ is the identity operator on $\mathcal{C}(G/H, \tau)$.*

Actually, this gives an expression of every element in $\mathcal{C}(G/H, \tau)$ as a wave packet of normalized Eisenstein integrals. The proof goes as follows. If $\mathcal{P}_\tau$ was not the identity, using Theorem 1 on the temperedness of the spectrum, one could find a non zero element of $\mathcal{A}_{temp}(G/H, \tau)$ which is orthogonal to the image of $\mathcal{P}_\tau$. Then, generalizing Theorem 5 to the truncated inner product of an Eisenstein integral with a general element of $\mathcal{A}_{temp}(G/H, \tau)$, this orthogonality can be explicitly described (cf. the evaluation of I before Theorem 9). As a result, this function has to be zero, a contradiction which proves the theorem.

The theorem translates to K -finite functions, involving representations and H -fixed distribution vectors.

The theorem can also be expressed with the unnormalized Eisenstein integrals. Then there are certain Plancherel factors involved. They are linked, as in the group case, to the intertwining integrals. Following the approach of A. Knapp and G. Zuckerman, [KZ], their computation is reduced to find an embedding of discrete series into principal series attached to minimal parabolic subgroups. For connected groups, this has been done by J. Carmona [C2].

8. Applications and open problems

Schwartz space for the hypergeometric Fourier transform

The image of a natural Schwartz space by the hypergeometric Fourier transform is characterized [D5]. The work uses the Plancherel formula of E. Opdam [Op], and the techniques mentioned above : theory of the constant term, C -functions, truncation ...

Generalized Schur orthogonality relations

Using the Plancherel formula for reductive symmetric spaces groups, K. Ankabout, [An], has proved generalized Schur orthogonality relations for generalized coefficients related to real reductive symmetric spaces. In particular, at least if we assume multiplicity one in the Plancherel formula, it implies the following:

There exists an explicit positive function d on G/H , such that, for almost all representations π occurring in the Plancherel formula, for ξ an H -fixed distribution vector of π occurring in the decomposition of the Dirac measure, there exists an

explicit non zero constant C_π such that, for all v, v' , K -finite vectors in the space of π :

$$\lim_{\varepsilon \rightarrow 0^+} \varepsilon^{n_\pi} \int_{G/H} e^{-\varepsilon d(x)} \langle \pi'(g)\xi, v \rangle \overline{\langle \pi'(g)\xi, v' \rangle} dx = C_\pi(v, v').$$

Here n_π is the dimension of the support of the Plancherel decomposition, around π . This refines and generalizes a work of Mirodikawa. It suggests to look for such type of relations in other situations.

$\mathbb{D}(G/H)$ -finite τ -spherical functions on reductive symmetric spaces

S. Souaifi [So] showed how these functions appear as linear combinations of derivatives along the complex parameter λ , of Eisenstein integrals. For K -finite functions, filtrations are introduced, whose subquotients are described in terms of induced representations. The starting point is an adaptation of ideas used by J. Franke to study spaces of automorphic forms. The use of the spectral decomposition by Langlands is replaced here by the use of the Plancherel formula. For reductive p -adic groups, and for the group case, I got similar results.

Invariant harmonic analysis on real reductive symmetric spaces

The goal is to study the H -invariant eigendistributions under $\mathbb{D}(G/H)$ on G/H and to express invariant measures on certain H -orbits in terms of these distributions (cf [D3] for the work of A.Bouaziz and P.Harinck for the group case and $G(\mathbb{C})/G(\mathbb{R})$, see also [OSe]).

Harmonic analysis on p -adic reductive symmetric spaces

For the group case, the Problems (b) and (c) of the Introduction have been solved by Harish-Chandra, up to the explicit description of the discrete series. In general, the problems are open (see [HH] for interesting structural results).

References

- [An] K.Ankabout, Relations d'orthogonalité de Schur généralisées pour les espaces symétriques réductifs , *J. Funct. Anal.*, 185 (2001), 63–110.
- [A] J. Arthur, A local trace formula, *Pub. Math. I.H.E.S* , 73 (1991), 5–96.
- [B1] E.van den Ban, The principal series for a reductive symmetric space I, *Ann. Sc. Ec. Norm. Sup.*, 21 (1988), 359–412.
- [B2] E.van den Ban, The principal series for a reductive symmetric space II, *J. of Funct. Analysis*, 109 (1992), 331–441.
- [BCD] E.van den Ban, J.Carmona, P.Delorme, Paquets d'ondes dans l'espace de Schwartz d'un espace symétrique réductif , *J. of Funct. Analysis* , 139 (1996), 225–243.
- [BS1] E.van den Ban , H.Schlichtkrull, Fourier transform on a semisimple symmetric space, Preprint Universiteit Utrecht, No 888, Nov. 1994.
- [BS2] E. van den Ban , H. Schlichtkrull, The most continuous part of the Plancherel decomposition for a reductive symmetric space , *Ann. of Math.*, 145 (1997) , 267–364.

- [BS3] E.van den Ban , H.Schlichtkrull, Harmonic analysis on reductive symmetric spaces , *European Congress of Mathematics, Barcelona, 2000, Vol 1*, 565–582, Birkhauser Verlag, Basel, 2001.
- [BS4] E.van den Ban , H.Schlichtkrull, The Plancherel decomposition for a reductive symmetric space I, II, Preprints 2001.
- [Be] J.N. Bernstein, On the support of the Plancherel measure, *J. Geom. Phys.*, 5 (1988), 663–710.
- [BrD] J.L.Brylinski, P.Delorme, Vecteurs distributions H -invariants pour les séries principales généralisées d’espaces symétriques réductifs et prolongement méromorphe d’intégrales d’Eisenstein , *Inv. Math.*, 109 (1992), 619–664.
- [C1] J.Carmona, it Terme constant des fonctions tempérées sur un espace symétrique réductif,, *J. fur Reine Angew. Math.*, 491 (1997), 17–63.
- [C2] J.Carmona, Plongement de séries discrètes pour un espace symétrique réductif, *J. of Funct. Analysis*, 182 (2001), 16–51.
- [CD1] J.Carmona, P.Delorme, Base méromorphe de vecteurs distributions H -invariants pour les séries principales généralisées d’espaces symétriques réductifs, *J. of Funct. Analysis*, 122 (1994), 152–221.
- [CD2] J.Carmona, P.Delorme, Transformation de Fourier pour les espaces symétriques réductifs , *Invent. Math.*, 134 (1998).
- [D1] P.Delorme, Intégrales d’Eisenstein pour les espaces symétriques réductifs. Tempérance. Majorations. Petite matrice B , *J. of Funct. Analysis*, 136 (1996), 422–509.
- [D2] P.Delorme, Troncature pour les espaces symétriques réductifs , *Acta Math.*, 179 (1997), 41–77.
- [D3] P.Delorme, Inversion des intégrales orbitales sur certains espaces symétriques réductifs, d’après A. Bouaziz et P. Harinck, *Séminaire Bourbaki, Vol. 1995-1996*, Astérisque 241, Exp. 810, 417–452.
- [D4] P.Delorme, Formule de Plancherel pour les espaces symétriques réductifs , *Ann. of Math.*, 147 (1998), 417–452.
- [D5] P.Delorme, Espace de Schwartz pour la transformation de Fourier hypergéométrique , Appendice par M. TINFOU, *J. Funct. Anal.* , 168 (1999), 239–312.
- [D6] P.Delorme, The Plancherel formula on reductive symmetric spaces from the point of view of the Schwartz space. Lectures for the European Summer School of Group Theory, CIRM, Luminy 2001.
- [F-J] M.Flensted-Jensen, Discrete series for semisimple symmetric spaces , *Ann. of Math.*, 111(1980), 253–311.
- [Ha] P.Harinck, Fonctions orbitales sur $G_{\mathbb{C}}/G_{\mathbb{R}}$. Formule d’inversion des intégrales orbitales et formule de Plancherel , *J. Funct. Anal.*, 153 (1998), 52–107.
- [H-C1] Harish-Chandra, Harmonic analysis on real reductive groups I , *J. Funct. Anal.*, 36 (1976), 1–35.
- [H-C2] Harish-Chandra, Harmonic analysis on real reductive groups II, *Inventiones Math.* , 36 (1976), 1–35.

- [H-C3] Harish-Chandra, Harmonic analysis on real reductive groups III. The Maass-Selberg relations and the Plancherel formula , *Ann. of Math.*, 104 (1976), 117–201.
- [He] S. Helgason, *Groups and geometric analysis*, Academic Press, 1984.
- [HH] A.G. Helminck, G.F. Helminck , A class of parabolic k -subgroups associated with symmetric k -varieties , *Trans. Amer. Math. Soc.*, 350 (1998), 4669–4684.
- [KZ] A. Knapp, G. Zuckerman, Classification of irreducible tempered representations of semisimple groups I , *Ann. of Math.* 116 (1982), 389–455.
- [Ol] G. Olafsson, Fourier and Poisson transformation associated to a semisimple symmetric space, *Inv. Math.*, 90 (1987), 1–51.
- [Op] E. Opdam, Cuspidal hypergeometric functions , dedicated to R. Askey, Part I, *Methods Appl. Anal.* 6 (1999) 67–80.
- [O1] T. Oshima, Fourier analysis on semisimple symmetric spaces, *Non commutative harmonic analysis and Lie groups (Marseille-Luminy, 1980)*, Lecture Notes in Math., vol 880, Springer, 1991, 357–369.
- [O2] T. Oshima, Asymptotic behaviour of spherical functions on semisimple symmetric spaces , *Adv. Studies in Pure Math.* 14 (1988), 561–601.
- [OM] T. Oshima, T. Matsuki, A description of discrete series for semisimple symmetric spaces, *Adv. Studies in Pure Math.*, 4 (1984), 331–390.
- [OSe] T. Oshima, J. Sekiguchi, Eigenspaces of invariant differential operators on a semisimple symmetric space, *Inv. Math.* 57 (1980), 1–81.
- [S] H. Schlichtkrull, The Langlands parameter of Flensted-Jensen’s discrete series for semisimple symmetric spaces , *J. of Funct Anal.*, 50 (1983), 133–150.
- [So] S. Souaifi , Fonctions $\mathbb{D}(G/H)$ -finies sur un espace symétrique réductif, Preprint, 2001, to appear in *J. of Funct Anal.*
- [V] D. Vogan, Irreducibility of discrete series representation for semisimple symmetric spaces , *Adv. Studies in Pure Math.*, 14 (1988), 191–221.
- [W] N. Wallach, *Real Reductive Groups II*, Academic Press, Inc., Boston, 1992.